

Higher National School of Hydraulic

The Library

Digital Repository of ENSH



المدرسة الوطنية العليا للري

المكتبة

المستودع الرقمي للمدرسة العليا للري



The title (العنوان):

Identification of groundwater potential zones using remote sensing, GIS, machine learning and electrical resistivity tomography techniques in Guelma basin, northeastern Algeria

The paper document Shelf mark P22-19 (: paper version not available)

APA Citation (توثيق APA):

Braham Massinissa, Boufekane Abdelmadjid, Bourenane Hamid (2022). *Identification of groundwater potential zones using remote sensing, GIS, machine learning and electrical resistivity tomography techniques in Guelma basin, northeastern Algeria.*

Geocarto International , vol 16(n°61), p 1-31 . DOI ou URL :

<https://www.tandfonline.com/doi/full/10.1080/10106049.2022.2063408?scroll=top&need>

Access=true&role=tab

The digital repository of the Higher National School for Hydraulics "Digital Repository of ENSH" is a platform for valuing the scientific production of the school's teachers and researchers.

Digital Repository of ENSH aims to limit scientific production, whether published or unpublished (theses, pedagogical publications, periodical articles, books...) and broadcasting it online.

Digital Repository of ENSH is built on the open DSpace software platform and is managed by the Library of the National Higher School for Hydraulics. <http://dspace.ensh.dz/jspui/>

المستودع الرقمي للمدرسة الوطنية العليا للري هو منصة خاصة بتمكين لإنتاج لأساتذة باحثي المدرسة.

يهدف المستودع الرقمي للمدرسة إلى حصر الإنتاج العلمي سواء كان منشورا أو غير منشور (طروحات، مطبوعات، مذكرات، مقالات، دوريات، كتب...) بثه على الخط.

المستودع الرقمي للمدرسة مبني على المنصة المفتوحة DSpace و يتم إدارته من طرف مديرية المكتبة للمدرسة العليا .

كل الحقوق محفوظة للمدرسة الوطنية العليا للري.

Abstract : In this research we assess and map groundwater potential in the Guelma Basin (northeastern Algeria) using an approach combining remote sensing, GIS, statistical and machine learning models. Four models were used including the frequency ratio model with both conventional (CFR) and modified (MFR) versions, the decision tree (DT), and the random forest (RF). For this purpose, firstly, thirteen hydro-geo-morphological variables influencing groundwater potential have been mapped using GIS and remote sensing techniques including elevation, slope, aspect, topographic wetness index, slope Length and Steepness factor, profile curvature, plan curvature, drainage density, distance to river, lineament and fault density, distance to faults and lineaments, lithology, and land use/land cover. Secondly, the groundwater potential was assessed and mapped based on the four models using the training data. Finally, the obtained groundwater potential maps of the four models have been validated using two approaches: (i) a statistical approach based on the receiver operating characteristics curves (ROC); (ii) a geophysical approach by interpreting the electrical resistivity tomography (ERT) results. The validation process gives the Random Forest method as the most accurate. The obtained map by this model is the main finding of this research, where the very high groundwater potential class occupies 8.25%. It is located mostly in the Guelma plain centre and in the northern part of the study area. The used approach and the obtained results may serve for water resource managers to improve groundwater resource planning and to resolve regional scale issues in this area or elsewhere.

Key words : Machine learning ; Groundwater potential ; Remote sensing, GIS ; Electrical resistivity tomography

Available from: - <https://www.tandfonline.com/doi/abs/10.1080/10106049.2022.2063408>